

Q P
623
.L33
1996

Address All Inquiries to the Publisher
Wiley-Liss, Inc., 605 Third Avenue, New York, NY 10158-0012

Copyright © 1996 Wiley-Liss, Inc.

Printed in the United States of America

The text of this book is printed on acid-free paper.

Under the conditions stated below the owner of copyright for this book hereby grants permission to users to make photocopy reproductions of any part or all of its contents for personal or internal organizational use, or for personal or internal use of specific clients. This consent is given on the condition that the copier pay the stated per-copy fee through the Copyright Clearance Center, Incorporated, 27 Congress Street, Salem, MA 01970, as listed in the most current issue of "Permissions to Photocopy" (Publisher's Fee List, distributed by CCC, Inc.), for copying beyond that permitted by sections 107 or 108 of the US Copyright Law. This consent does not extend to other kinds of copying, such as copying for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale.

While the authors, editors, and publisher believe that drug selection and dosage and the specification and usage of equipment and devices, as set forth in this book, are in accord with current recommendations and practice at the time of publication, they accept no legal responsibility for any errors or omissions, and make no warranty, express or implied, with respect to material contained herein. In view of ongoing research, equipment modifications, changes in governmental regulations and the constant flow of information relating to drug therapy, drug reactions, and the use of equipment and devices, the reader is urged to review and evaluate the information provided in the package insert or instructions for each drug, piece of equipment, or device for, among other things, any changes in the instructions or indication of dosage or usage and for added warnings and precautions.

Library of Congress Cataloging in Publication Data:

A laboratory guide to RNA : isolation, analysis, and synthesis /
edited by Paul A. Krieg.

p. cm.

Includes bibliographical references and index.

ISBN 0-471-12536-9 (paper : alk. paper)

1. RNA—Analysis—Laboratory manuals. I. Krieg, Paul A., 1953—

QP623.L33 1996

574.87'3283—dc20

96-19936

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

cze 8305

CON

D. CRAIG
7703 F

FRANCESCO
della R

MARGARET
Science

Yael Bibb
try, Heb
Israel

ERIC BUXY
92008

DANIELA C
Medicin

CARL G. C
ence Ce

KIRAN CHA
tistry of

ALEX CHEI
Way, Pa

LATA CHE
Dentist

ONDINE B
versity

RONALD A
Euclid

JEANINE D
and De

FRANÇOIS
CA 920

SUZANNE
Science

- chemical means to "freeze" the ribosomes engaged in translation at their sites on mRNAs. It should be pointed out, however, that incubation of the cells with cycloheximide for extended periods might lead to an artifactual loading of ribosomes onto mRNAs. Hence cycloheximide should be added to the culture 1–15 min prior to the cell harvest.
- f. The gradients should be cold on loading of the sample. They can be prepared at room temperature and then either rapidly cooled by placing them in ice for about 30 min, or kept in the cold room prior to the sample loading. Gradients are known to remain stable overnight.
 - g. Occasionally, the lysis solution is not clear, due to precipitation of the detergents. It can still be used, but should be mixed well before aliquoting.
 - h. Heparin is a potent RNase inhibitor, but at high concentrations it might lead to nuclear lysis. To avoid contamination of the cytoplasmic extract the concentration of heparin should not exceed 100 $\mu\text{g}/\text{mL}$ prior to the removal of nuclei. Heparin can be substituted, however, by RNA guard (Pharmacia) or RNasin (Promega) at a concentration of 150–300 units mL . These RNase inhibitors require a minimum of 1 mM DTT in the lysis buffer for their activity and should be added directly to the tissues or cells prior to homogenization.
 - i. SDS is an efficient anti-RNase detergent. However, all steps in which SDS is involved should be carried out at room temperature as this detergent precipitates out of solution at $<15^\circ\text{C}$.
 - j. Partitioning of the sucrose gradient into two fractions saves time and reagents, particularly if the only relevant parameter is the distribution of an mRNA between polysome and subpolysomes (Fig. 5.1b). However, due to the poor resolution of this approach, potentially important information will be lost. Analysis of the distribution of an mRNA along a gradient which is divided into 12 fractions will enable monitoring whether an mRNA is shifted from heavy to light polysomes and allow an estimate of the maximum number of ribosomes loaded on the mRNA (Figs. 5.1c, 5.1d, 5.2b, 5.2c, 5.3b, and 5.3c). The border between the subpolysomal fractions is indicated by the low point in the profile immediately following the polysomal peaks (Figs. 5.1a, 5.2a, and 5.3a).
 - k. RNA can be isolated by alternative methods as described below.
 1. If the polysomes are derived from a protein-rich source like rat liver, or oocytes and embryos of *Xenopus*, it is often helpful to digest the proteins using the proteinase K/SDS/phenol procedure (Probst et al., 1979). In this case, adjust the method as follows: (1) add 400 μL of PK buffer, instead of the 0.5% SDS buffer; (2) incubate for 30 min at room temperature; and (3) extract the RNA with an equal volume of chloroform as described in steps 30–38, with appropriate volume adjustments.
 - m. Ideally, RNA will be extracted from all gradient fractions with similar efficiency. This will permit a reliable assessment of the distribution of a specific mRNA among these fractions. However, the large amount of

prote
tion.
inter
phase
speed
remo
with
n. Mess
using
plasm
mal
polys
cDNA
In
SDS
poly-
um.
o. Both
active
addit

COMMENT

Examples

Translation

Cells. The regulated in lease from lymphosarcoma proliferate w This translat polysomes f (Fig. 5.1). another hous associated wit apparent dist evident whe somes and S However, do coding rpl3 this class of mRNAs is o populations e active (poly e By compari derived from